

## PROTEASES FROM GRAM-POSITIVE ORGANISMS

### FIELD OF THE INVENTION

The present invention relates to metallo-proteases derived from gram-positive microorganisms. The present invention provides nucleic acid and amino acid sequences of a metallo-protease identified in *Bacillus subtilis*. The present invention also provides methods for the production of the protease in host cells as well as the production of heterologous proteins in a host cell having a mutation or deletion of part or all of the proteases of the present invention.

### BACKGROUND OF THE INVENTION

Gram-positive microorganisms, such as members of the group *Bacillus*, have been used for large-scale industrial fermentation due, in part, to their ability to secrete their fermentation products into the culture media. In gram-positive bacteria, secreted proteins are exported across a cell membrane and a cell wall, and then are subsequently released into the external media usually maintaining their native conformation.

Various gram-positive microorganisms are known to secrete extracellular and/or intracellular protease at some stage in their life cycles. Many proteases are produced in large quantities for industrial purposes. A negative aspect of the presence of proteases in gram-positive organisms is their contribution to the overall degradation of secreted heterologous or foreign proteins.

The classification of proteases found in microorganisms is based on their catalytic mechanism which results in four groups: the serine proteases; metallo-proteases; cysteine proteases; and aspartic proteases. These categories, in general, can be distinguished by their sensitivity to various inhibitors. For example, the serine proteases are inhibited by phenylmethylsulfonylfluoride (PMSF) and diisopropylfluorophosphate (DIFP); the metallo-proteases by chelating agents; the cysteine enzymes by iodoacetamide and heavy metals and the aspartic proteases by pepstatin. The serine proteases have alkaline pH optima, the metalloproteases are optimally active around neutrality, and the cysteine and aspartic enzymes have acidic pH optima (*Biotechnology Handbooks, Bacillus*, vol. 2, edited by Harwood, 1989 Plenum Press, New York).

Metallo-proteases form the most diverse of the catalytic types of proteases. Family M23 contains bacterial enzymes such as the  $\beta$ -lytic endopeptidases of *Lysobacter* and

-- 2 --

*Achromobacter* and the *Pseudomonas* LasA protein and have specificity for Gly bonds, especially in Gly-Gly+Xaa-sequences (Methods in Enzymology, vol. 248, Academic Press, Inc. 1994). The enzymes of the M23 family contain zinc and a conserved His-Xaa-His motif.

### SUMMARY OF THE INVENTION

The present invention relates to the discovery of a heretofore unknown metallo-protease (MP) found in gram positive microorganisms, uses of the MP in industrial applications, and advantageous strain improvements based on genetically engineering such microorganisms to delete, underexpress or overexpress that MP. Due to the overall relatedness of MP with *Pseudomonas* lasA protein, including the presence of the motif His-Xaa-His, MP appears to be a member of the metallo-protease family M23.

Applicant's discovery, in addition to providing a new and useful protease and methods of detecting DNA encoding such proteases in a gram positive microorganism, provides several advantages which may facilitate optimization and/or modification of strains of gram positive microorganisms, such as *Bacillus*, for expression of desired, e.g. heterologous, proteins. Such optimizations, as described below in detail, allow the construction of strains having decreased proteolytic degradation of desired expression products.

Applicant's invention is further based on the discovery of the presence of MP's in Gram-positive microorganisms. The Gram-positive microorganism may be *Bacillus* and may also be selected from the group consisting of *Bacillus subtilis*, *Bacillus stearothermophilus*, *Bacillus licheniformis* and *Bacillus amyloliquifaciens*. The present invention further relies on the discovery that naturally occurring MP is encoded by nucleic acid found about 2248 kb from the point of origin of *Bacillus subtilis* I-168 strain (*Bacillus* Genetic Stock Center, accession number 1A1, Columbus, Ohio). The present invention relates to the MP encoded thereby, as well as the nucleic acid and amino acid molecules having the sequences disclosed in Figures 1A-1O.

The present invention thus provides methods for detecting gram positive microorganism homologs of *B. subtilis* MP that comprises hybridizing part or all of the nucleic acid encoding *B. subtilis* MP with nucleic acid derived from gram-positive organisms, either of genomic or cDNA origin. Accordingly, the present invention provides a method for detecting a gram-positive microorganism MP, comprising the steps of hybridizing gram-positive microorganism nucleic acid under low stringency conditions to a probe, wherein the probe

-- 3 --

comprises part or all of the nucleic acid sequence shown in Figures 1A-1O; and isolating gram-positive nucleic acid which hybridizes to said probe.

In a preferred embodiment, the *Bacillus* is selected from the group consisting of *B. licheniformis*, *B. lentus*, *B. brevis*, *B. stearothermophilus*, *B. alkalophilus*, *B. amyloliquefaciens*, *B. coagulans*, *B. circulans*, *B. lautus* and *B. thuringiensis*.

The production of desired heterologous proteins or polypeptides in gram-positive microorganisms may be hindered by the presence of one or more proteases, including MP, which degrade the produced heterologous protein or polypeptide. One advantage of the present invention is that it provides methods and expression systems which can be used to prevent that degradation, thereby enhancing yields of the desired heterologous protein or polypeptide. Accordingly, the present invention provides a gram-positive microorganism having a mutation or deletion of part or all of the gene encoding MP, which results in the inactivation of the MP proteolytic activity, either alone or in combination with mutations in other proteases, such as apr, npr, epr, mpr, bpf or isp for example, or other proteases known to those of skill in the art. In one embodiment of the present invention, the gram-positive organism is a member of the genus *Bacillus*. In another embodiment, the *Bacillus* is selected from the group consisting of *B. subtilis*, *B. licheniformis*, *B. lentus*, *B. brevis*, *B. stearothermophilus*, *B. alkalophilus*, *B. amyloliquefaciens*, *B. coagulans*, *B. circulans*, *B. lautus* and *Bacillus thuringiensis*. In a further preferred embodiment, the *Bacillus* is *Bacillus subtilis*.

In another aspect, the gram-positive host having one or more metallo-protease deletions or mutations is further genetically engineered to produce a desired protein. In one embodiment of the present invention, the desired protein is heterologous to the gram-positive host cell. In another embodiment, the desired protein is homologous to the host cell. The present invention encompasses a gram-positive host cell having a deletion, mutation or interruption of the nucleic acid encoding the naturally occurring homologous protein, such as a protease, and having nucleic acid encoding the homologous protein re-introduced in a recombinant form. In another embodiment, the host cell produces the homologous protein. Accordingly, the present invention also provides methods and expression systems for reducing degradation of heterologous proteins produced in gram-positive microorganisms. The gram-positive microorganism may be normally sporulating or non-sporulating. In a preferred embodiment, the gram positive host cell is a *Bacillus*. In another preferred embodiment, the *Bacillus* host cell is *Bacillus*. In another embodiment, the *Bacillus* is selected from the group

-- 4 --

consisting of *B. subtilis*, *B. licheniformis*, *B. lentus*, *B. brevis*, *B. stearothermophilus*, *B. alkalophilus*, *B. amyloliquefaciens*, *B. coagulans*, *B. circulans*, *B. lautus* and *Bacillus thuringiensis*.

Naturally occurring gram positive MP as well as proteolytically active amino acid variations or derivatives thereof, have application in the textile industry, in cleaning compositions and in animal feed. The metallo-protease MP may be used alone or in combination with other enzymes and/or mediators or enhancers. Accordingly, in a further aspect of the present invention, gram-positive MP is produced on an industrial fermentation scale in a microbial host expression system. The present invention provides a cleaning composition comprising a metalloprotease, MP, having the amino acid sequence shown in Figures 1A-1O or the amino acid encoded by the MP nucleic acid found at about 2248 kilobases from the point of origin of *Bacillus subtilis*. Also provided are cleaning compositions comprising a metalloprotease having at least 80%, at least 90%, or at least 95% homology with the amino acid sequence shown in Figures 1A-1O or comprising a metalloprotease encoded by a gene that hybridizes with the nucleic acid shown in Figures 1A-1O under high stringency conditions.

Further there is provided an animal feed comprising a metalloprotease, MP, having the amino acid sequence shown in Figures 1A-1O. Also provided are animal feeds comprising a metalloprotease having at least 80%, at least 90%, and at least 95% homology with the amino acid sequence shown in Figures 1A-1O or comprising a metalloprotease encoded by a gene that hybridizes with the nucleic acid shown in Figures 1A-1O under high stringency conditions.

Also provided is a composition for the treatment of a textile comprising a metalloprotease, MP, having the amino acid sequence shown in Figures 1A-1O. Also provided are compositions for the treatment of a textile comprising a metalloprotease having at least 80%, at least 90%, or at least 95% homology with the amino acid sequence shown in Figures 1A-1O or comprising a metalloprotease encoded by a gene that hybridizes with the nucleic acid shown in Figures 1A-1O under high stringency conditions.

### **BRIEF DESCRIPTION OF THE DRAWINGS**

Figures 1A-1O shows the DNA and amino acid sequence for *Bacillus subtilis* MP.

Figure 2 show an amino acid alignment of *Bacillus subtilis* MP (designated as YOMI) and *Pseudomonas* LasA. The amino acid motif H-X-H is noted at amino acid 308 -310 in LasA.

## DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

### Definitions

As used herein, the genus *Bacillus* includes all members known to those of skill in the art, including but not limited to *B. subtilis*, *B. licheniformis*, *B. lentus*, *B. brevis*, *B. stearothermophilus*, *B. alkalophilus*, *B. amyloliquefaciens*, *B. coagulans*, *B. ciculans*, *B. lautus* and *B. thuringiensis*.

The present invention relates to a newly characterized metallo-protease (MP) from gram positive organisms. In a preferred embodiment, the metallo-protease is obtainable from a gram-positive organism which is a *Bacillus*. In another preferred embodiment, the metallo-protease is obtainable from a *Bacillus* which is selected from the group consisting of *B. subtilis*, *B. licheniformis*, *B. lentus*, *B. brevis*, *B. stearothermophilus*, *B. alkalophilus*, *B. amyloliquefaciens*, *B. coagulans*, *B. ciculans*, *B. lautus* and *B. thuringiensis*.

In another preferred embodiment, the gram-positive organism is *Bacillus subtilis* and MP has the amino acid sequence encoded by the nucleic acid molecule having the sequence that occurs around 2248 kilobases from the point of origin of *Bacillus subtilis* I-168.

In another preferred embodiment, *Bacillus subtilis* has the nucleic acid and amino acid sequence as shown in Figures 1A-1O. The present invention encompasses the use of amino acid variations of the amino acid sequences disclosed in Figures 1A-1O that have proteolytic activity. Such proteolytic amino acid variants can be used in the textile industry, animal feed and in cleaning compositions. The present invention also encompasses the use of *B. subtilis* amino acid variations or derivatives that are not proteolytically active. DNA encoding such variants can be used in methods designed to delete or mutate the naturally occurring host cell MP.

As used herein, "nucleic acid" refers to a nucleotide or polynucleotide sequence, and fragments or portions thereof, and to DNA or RNA of genomic or synthetic origin which may be double-stranded or single-stranded, whether representing the sense or antisense strand. As used herein "amino acid" refers to peptide or protein sequences or portions thereof. A "polynucleotide homolog" as used herein refers to a gram-positive microorganism polynucleotide that has at least 80%, at least 90% and at least 95% identity to *B. subtilis* MP, or which is capable of hybridizing to *B. subtilis* MP under conditions of high stringency and which encodes an amino acid sequence having metallo-protease activity.

-- 6 --

The terms "isolated" or "purified" as used herein refer to a nucleic acid or amino acid that is removed from at least one component with which it is naturally associated.

As used herein, the term "heterologous protein" refers to a protein or polypeptide that does not naturally occur in a gram-positive host cell. Examples of heterologous proteins include enzymes such as hydrolases including proteases, cellulases, amylases, carbohydrases, and lipases; isomerases such as racemases, epimerases, tautomerases, or mutases; transferases, kinases and phosphatases. The heterologous gene may encode therapeutically significant proteins or peptides, such as growth factors, cytokines, ligands, receptors and inhibitors, as well as vaccines and antibodies. The gene may encode commercially important industrial proteins or peptides, such as proteases, carbohydrases such as amylases and glucoamylases, cellulases, oxidases and lipases. The gene of interest may be a naturally occurring gene, a mutated gene or a synthetic gene.

The term "homologous protein" refers to a protein or polypeptide native or naturally occurring in a gram-positive host cell. The invention includes host cells producing the homologous protein via recombinant DNA technology. The present invention encompasses a gram-positive host cell having a deletion or interruption of the nucleic acid encoding the naturally occurring homologous protein, such as a protease, and having nucleic acid encoding the homologous protein re-introduced in a recombinant form. In another embodiment, the host cell produces the homologous protein.

As used herein, the term "overexpressing" when referring to the production of a protein in a host cell means that the protein is produced in greater amounts than its production in its naturally occurring environment.

As used herein, the phrase "proteolytic activity" refers to a protein that is able to hydrolyze a peptide bond. Enzymes having proteolytic activity are described in Enzyme Nomenclature, 1992, edited Webb Academic Press, Inc.

#### **DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS**

The unexpected discovery of the metallo-protease M23 family member, designated herein as MP, found in translated, uncharacterized *B.subtilis* genomic sequences provides a basis for producing host cells, expression methods and systems which can be used to prevent the degradation of recombinantly produced heterologous proteins.

Accordingly, in a preferred embodiment, the host cell is a gram-positive host cell that has a deletion or mutation in the naturally occurring nucleic acid encoding MP said mutation

-- 7 --

resulting in deletion or inactivation of the production by the host cell of the MP proteolytic gene product. The host cell may additionally be genetically engineered to produce a desired protein or polypeptide.

It may also be desired to genetically engineer host cells of any type to produce a gram-positive MP. Such host cells are used in large scale fermentation to produce large quantities of the protease which may be isolated or purified and used in cleaning products, such as detergents, in textile treatments and as animal feed additives.

### I. MP Sequences

The nucleic acid sequence and amino acid sequence for *Bacillus subtilis* MP are shown in Figures 1A-1O. As will be understood by the skilled artisan, due to the degeneracy of the genetic code, a variety of polynucleotides can encode the *Bacillus subtilis* MP having the amino acid sequence shown in Figures 1A-1O. The present invention encompasses all such polynucleotides.

The present invention encompasses the use of MP polynucleotide homologs encoding gram-positive microorganism MPs which have at least 80%, or at least 90% or at least 95% identity to *B. subtilis* MP shown in Figures 1A-1O as long as the homolog encodes a protein that has proteolytic activity.

Gram-positive polynucleotide homologs of *B. subtilis* MP may be obtained by standard procedures known in the art from, for example, cloned DNA (*e.g.*, a DNA "library"), genomic DNA libraries, by chemical synthesis once identified, by cDNA cloning, or by the cloning of genomic DNA, or fragments thereof, purified from a desired cell. (*See*, for example, Sambrook *et al.*, 1989, Molecular Cloning, A Laboratory Manual, 2d Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York; Glover, D.M. (ed.), 1985, DNA Cloning: A Practical Approach, MRL Press, Ltd., Oxford, U.K. Vol. I, II.) A preferred source is from genomic DNA.

As will be understood by those of skill in the art, the polynucleotide sequence disclosed in Figures 1A-1O may reflect inadvertent errors inherent to nucleic acid sequencing technology. Moreover, the sequence of polynucleotides derived from related species, *e.g.*, other *Bacillus*, will contain variations to the sequences specifically disclosed herein. Nonetheless, one of ordinary skill in the art is fully capable of determining the correct sequences from the information provided herein regarding the invention. For example, as described below, it is possible to identify the MP of the invention by virtue of its location in the microorganism's genome. The present invention

-- 8 --

encompasses the naturally occurring nucleic acid molecule having the nucleic acid sequence obtained from the genomic sequence of *Bacillus* species.

Nucleic acid encoding *Bacillus subtilis* MP starts around 2248 kilobases counting from the point of origin in the *Bacillus subtilis* strain I-168 (Anagnostopala, 1961, J. Bacteriol. 81:741-746 or *Bacillus* Genomic Stock Center, accession 1A1, Columbus, Ohio). The *Bacillus subtilis* point of origin has been described in Ogasawara, N. (1995, Microbiology 141:Pt.2 257-59). *Bacillus subtilis* MP has a length of 2285 amino acids. Based upon the location of the DNA encoding *Bacillus subtilis* MP, naturally occurring *B. subtilis* MP can be obtained by methods known to those of skill in the art including PCR technology.

Oligonucleotide sequences or primers of about 10-30 nucleotides in length can be designed from the polynucleotide sequence disclosed in Figures 1A-1O and used in PCR technology to isolate the naturally occurring sequence from *B. subtilis* genomic sequences.

Another general strategy for the "cloning" of *B. subtilis* genomic DNA pieces for sequencing uses inverse PCR. A known region is scanned for a set of appropriate restriction enzyme cleavage sites and inverse PCR is performed with a set of DNA primers determined from the outermost DNA sequence. The DNA fragments from the inverse PCR are directly used as template in the sequencing reaction. The newly derived sequences can be used to design new oligonucleotides. These new oligonucleotides are used to amplify DNA fragments with genomic DNA as template. The sequence determination on both strands of a DNA region is finished by applying a primer walking strategy on the genomic PCR fragments. The benefit of multiple starting points in the primer walking results from the series of inverse PCR fragments with different sizes of new "cloned" DNA pieces. From the most external DNA sequence a new round of inverse PCR is started. The whole inverse PCR strategy is based on the sequential use of conventional taq polymerase and the use of long range inverse PCR in those cases in which the taq polymerase failed to amplify DNA fragments. Nucleic acid sequencing is performed using standard technology. One method for nucleic acid sequencing involves the use of a Perkin-Elmer Applied Biosystems 373 DNA sequencer (Perkin-Elmer, Foster City, California).

Nucleic acid sequences derived from genomic DNA may contain regulatory regions in addition to coding regions. Whatever the source, the isolated MP gene should be molecularly cloned into a suitable vector for propagation of the gene.

In the molecular cloning of the gene from genomic DNA, DNA fragments are generated, some of which will encode the desired gene. The DNA may be cleaved at specific sites using various restriction enzymes. Alternatively, one may use DNase in the presence of manganese to



-- 9 --

fragment the DNA, or the DNA can be physically sheared, as for example, by sonication. The linear DNA fragments can then be separated according to size by standard techniques, including but not limited to, agarose and polyacrylamide gel electrophoresis and column chromatography.

Once the DNA fragments are generated, identification of the specific DNA fragment containing the MP may be accomplished in a number of ways. For example, a *B.subtilis* MP gene of the present invention or its specific RNA, or a fragment thereof, such as a probe or primer, may be isolated and labeled and then used in hybridization assays to detect a gram-positive MP gene. (Benton, W. and Davis, R., 1977, Science 196:180; Grunstein, M. And Hogness, D., 1975, Proc. Natl. Acad. Sci. USA 72:3961). Those DNA fragments sharing substantial sequence similarity to the probe will hybridize under stringent conditions.

Accordingly, the present invention provides a method for the detection of gram-positive MP polynucleotide homologs which comprises hybridizing part or all of a nucleic acid sequence of *B. subtilis* MP with gram-positive microorganism nucleic acid of either genomic or cDNA origin.

Also included within the scope of the present invention is the use of gram-positive microorganism polynucleotide sequences that are capable of hybridizing to the nucleotide sequence of *B.subtilis* MP under conditions of intermediate to maximal stringency. Hybridization conditions are based on the melting temperature ( $T_m$ ) of the nucleic acid binding complex, as taught in Berger and Kimmel (1987, Guide to Molecular Cloning Techniques, Methods in Enzymology, Vol 152, Academic Press, San Diego CA) incorporated herein by reference, and confer a defined "stringency" as explained below.

"Maximum stringency" typically occurs at about  $T_m - 5^\circ\text{C}$  ( $5^\circ\text{C}$  below the  $T_m$  of the probe); "high stringency" at about  $5^\circ\text{C}$  to  $10^\circ\text{C}$  below  $T_m$ ; "intermediate stringency" at about  $10^\circ\text{C}$  to  $20^\circ\text{C}$  below  $T_m$ ; and "low stringency" at about  $20^\circ\text{C}$  to  $25^\circ\text{C}$  below  $T_m$ . As will be understood by those of skill in the art, a maximum stringency hybridization can be used to identify or detect identical polynucleotide sequences while an intermediate or low stringency hybridization can be used to identify or detect polynucleotide sequence homologs.

The term "hybridization" as used herein shall include "the process by which a strand of nucleic acid joins with a complementary strand through base pairing" (Coombs J (1994) Dictionary of Biotechnology, Stockton Press, New York NY).

The process of amplification as carried out in polymerase chain reaction (PCR) technologies is described in Dieffenbach CW and GS Dveksler (1995, PCR Primer, a

-- 10 --

Laboratory Manual, Cold Spring Harbor Press, Plainview NY). A nucleic acid sequence of at least about 10 nucleotides and as many as about 60 nucleotides from *B. subtilis* MP preferably about 12 to 30 nucleotides, and more preferably about 20-25 nucleotides can be used as a probe or PCR primer.

5 The *B. subtilis* MP amino acid sequences (shown in Figures 1A-1O) were identified via a BLAST search (Altschul, Stephen, Basic local alignment search tool, J. Mol. Biol. 215:403-410) of *Bacillus subtilis* genomic nucleic acid sequences. *B. subtilis* MP (YOMI) was identified by its overall nucleic acid identity to the metallo-protease, *Pseudomonas lasA*, including the presence of the catalytic domain H-X-H as shown in Figure 2.

## 10 II. Expression Systems

The present invention provides host cells, expression methods and systems for the enhanced production and secretion of desired heterologous or homologous proteins in gram-positive microorganisms. In one embodiment, a host cell is genetically engineered to have a deletion or mutation in the gene encoding a gram-positive MP such that the respective activity is deleted. In another embodiment of the present invention, a gram-positive microorganism is genetically engineered to produce a metallo-protease of the present invention.

### 15 Inactivation of a gram-positive metallo-protease in a host cell

20 Producing an expression host cell incapable of producing the naturally occurring metallo-protease necessitates the replacement and/or inactivation of the naturally occurring gene from the genome of the host cell. In a preferred embodiment, the mutation is a non-reverting mutation.

25 One method for mutating nucleic acid encoding a gram-positive metallo-protease is to clone the nucleic acid or part thereof, modify the nucleic acid by site directed mutagenesis and reintroduce the mutated nucleic acid into the cell on a plasmid. By homologous recombination, the mutated gene may be introduced into the chromosome. In the parent host cell, the result is that the naturally occurring nucleic acid and the mutated nucleic acid are located in tandem on the chromosome. After a second recombination, the modified sequence is left in the  
30 chromosome having thereby effectively introduced the mutation into the chromosomal gene for progeny of the parent host cell.

Another method for inactivating the metallo-protease proteolytic activity is through deleting the chromosomal gene copy. In a preferred embodiment, the entire gene is deleted,

-- 11 --

the deletion occurring in such a way as to make reversion impossible. In another preferred embodiment, a partial deletion is produced, provided that the nucleic acid sequence left in the chromosome is too short for homologous recombination with a plasmid encoded metallo-protease gene. In another preferred embodiment, nucleic acid encoding the catalytic amino acid residues are deleted.

Deletion of the naturally occurring gram-positive microorganism metallo-protease can be carried out as follows. A metallo-protease gene including its 5' and 3' regions is isolated and inserted into a cloning vector. The coding region of the metallo-protease gene is deleted from the vector *in vitro*, leaving behind a sufficient amount of the 5' and 3' flanking sequences to provide for homologous recombination with the naturally occurring gene in the parent host cell. The vector is then transformed into the gram-positive host cell. The vector integrates into the chromosome via homologous recombination in the flanking regions. This method leads to a gram-positive strain in which the protease gene has been deleted.

The vector used in an integration method is preferably a plasmid. A selectable marker may be included to allow for ease of identification of desired recombinant microorganisms. Additionally, as will be appreciated by one of skill in the art, the vector is preferably one which can be selectively integrated into the chromosome. This can be achieved by introducing an inducible origin of replication, for example, a temperature sensitive origin into the plasmid. By growing the transformants at a temperature to which the origin of replication is sensitive, the replication function of the plasmid is inactivated, thereby providing a means for selection of chromosomal integrants. Integrants may be selected for growth at high temperatures in the presence of the selectable marker, such as an antibiotic. Integration mechanisms are described in WO 88/06623.

Integration by the Campbell-type mechanism can take place in the 5' flanking region of the protease gene, resulting in a protease positive strain carrying the entire plasmid vector in the chromosome in the metallo-protease locus. Since illegitimate recombination will give different results it will be necessary to determine whether the complete gene has been deleted, such as through nucleic acid sequencing or restriction maps.

Another method of inactivating the naturally occurring metallo-protease gene is to mutagenize the chromosomal gene copy by transforming a gram-positive microorganism with oligonucleotides which are mutagenic. Alternatively, the chromosomal metallo-protease gene can be replaced with a mutant gene by homologous recombination.

-- 12 --

The present invention encompasses host cells having additional protease deletions or mutations, such as deletions or mutations in apr, npr, epr, mpr and others known to those of skill in the art.

One assay for the detection of mutants involves growing the *Bacillus* host cell on medium containing a protease substrate and measuring the appearance or lack thereof, of a zone of clearing or halo around the colonies. Host cells which have an inactive protease will exhibit little or no halo around the colonies.

### III. Production of Metallo-protease

For production of metallo-protease in a host cell, an expression vector comprising at least one copy of nucleic acid encoding a gram-positive microorganism MP, and preferably comprising multiple copies, is transformed into the host cell under conditions suitable for expression of the metallo-protease. In accordance with the present invention, polynucleotides which encode a gram-positive microorganism MP, or fragments thereof, or fusion proteins or polynucleotide homolog sequences that encode amino acid variants of *B. subtilis* MP, may be used to generate recombinant DNA molecules that direct their expression in host cells. In a preferred embodiment, the gram-positive host cell belongs to the genus *Bacillus*. In another preferred embodiment, the gram positive host cell is *B. subtilis*.

As will be understood by those of skill in the art, it may be advantageous to produce polynucleotide sequences possessing non-naturally occurring codons. Codons preferred by a particular gram-positive host cell (Murray E et al (1989) Nuc Acids Res 17:477-508) can be selected, for example, to increase the rate of expression or to produce recombinant RNA transcripts having desirable properties, such as a longer half-life, than transcripts produced from naturally occurring sequence.

Altered MP polynucleotide sequences which may be used in accordance with the invention include deletions, insertions or substitutions of different nucleotide residues resulting in a polynucleotide that encodes the same or a functionally equivalent MP homolog, respectively. As used herein a "deletion" is defined as a change in either nucleotide or amino acid sequence in which one or more nucleotides or amino acid residues, respectively, are absent.

As used herein an "insertion" or "addition" is that change in a nucleotide or amino acid sequence which has resulted in the addition of one or more nucleotides or amino acid residues, respectively, as compared to the naturally occurring MP.

-- 13 --

As used herein "substitution" results from the replacement of one or more nucleotides or amino acids by different nucleotides or amino acids, respectively.

The encoded protein may also show deletions, insertions or substitutions of amino acid residues which produce a silent change and result in a functionally equivalent MP variant.

5 Deliberate amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues as long as the variant retains the ability to modulate secretion. For example, negatively charged amino acids include aspartic acid and glutamic acid; positively charged amino acids include lysine and arginine; and amino acids with uncharged polar head groups having similar hydrophilicity  
10 values include leucine, isoleucine, valine; glycine, alanine; asparagine, glutamine; serine, threonine, phenylalanine, and tyrosine.

The MP polynucleotides of the present invention may be engineered in order to modify the cloning, processing and/or expression of the gene product. For example, mutations may be introduced using techniques which are well known in the art, eg, site-directed mutagenesis to  
15 insert new restriction sites, to alter glycosylation patterns or to change codon preference, for example.

In one embodiment of the present invention, a gram-positive microorganism MP polynucleotide may be ligated to a heterologous sequence to encode a fusion protein. A fusion protein may also be engineered to contain a cleavage site located between the metallo-protease  
20 nucleotide sequence and the heterologous protein sequence, so that the metallo-protease may be cleaved and purified away from the heterologous moiety.

#### IV. Vector Sequences

Expression vectors used in expressing the metallo-proteases of the present invention in  
25 gram-positive microorganisms comprise at least one promoter associated with a metallo-protease selected from the group consisting of MP, which promoter is functional in the host cell. In one embodiment of the present invention, the promoter is the wild-type promoter for the selected metallo-protease and in another embodiment of the present invention, the promoter is heterologous to the metallo-protease, but still functional in the host cell. In one  
30 preferred embodiment of the present invention, nucleic acid encoding the metallo-protease is stably integrated into the microorganism genome.

In a preferred embodiment, the expression vector contains a multiple cloning site cassette which preferably comprises at least one restriction endonuclease site unique to the

-- 14 --

vector, to facilitate ease of nucleic acid manipulation. In a preferred embodiment, the vector also comprises one or more selectable markers. As used herein, the term selectable marker refers to a gene capable of expression in the gram-positive host which allows for ease of selection of those hosts containing the vector. Examples of such selectable markers include but are not limited to antibiotics, such as, erythromycin, actinomycin, chloramphenicol and tetracycline.

#### V. Transformation

A variety of host cells can be used for the production *Bacillus subtilis* MP or MP homologs including bacterial, fungal, mammalian and insects cells. General transformation procedures are taught in Current Protocols In Molecular Biology (vol. 1, edited by Ausubel et al., John Wiley & Sons, Inc. 1987, Chapter 9) and include calcium phosphate methods, transformation using DEAE-Dextran and electroporation. Plant transformation methods are taught in Rodriguez (WO 95/14099, published 26 May 1995).

In a preferred embodiment, the host cell is a gram-positive microorganism and in another preferred embodiment, the host cell is *Bacillus*. In one embodiment of the present invention, nucleic acid encoding one or more MP(s) of the present invention is introduced into a host cell via an expression vector capable of replicating within the *Bacillus* host cell. Suitable replicating plasmids for *Bacillus* are described in Molecular Biological Methods for *Bacillus*, Ed. Harwood and Cutting, John Wiley & Sons, 1990, hereby expressly incorporated by reference; see chapter 3 on plasmids. Suitable replicating plasmids for *B. subtilis* are listed on page 92.

In another embodiment, where it is desired to produce the MP for use in cleaning compositions, nucleic acid encoding MP is stably integrated into the microorganism genome. Preferred host cells are gram-positive host cells. Another preferred host is *Bacillus*. Another preferred host is *Bacillus subtilis*. Several strategies have been described in the literature for the direct cloning of DNA in *Bacillus*. Plasmid marker rescue transformation involves the uptake of a donor plasmid by competent cells carrying a partially homologous resident plasmid (Contente et al., Plasmid 2:555-571 (1979); Haima et al., Mol. Gen. Genet. 223:185-191 (1990); Weinrauch et al., J. Bacteriol. 154(3):1077-1087 (1983); and Weinrauch et al., J. Bacteriol. 169(3):1205-1211 (1987)). The incoming donor plasmid recombines with the homologous region of the resident "helper" plasmid in a process that mimics chromosomal transformation.

-- 15 --

Protoplast transformation is described for *B. subtilis* in Chang and Cohen, (1979) Mol. Gen. Genet 168:111-115; for *B. megaterium* in Vorobjeva et al., (1980) FEMS Microbiol. Letters 7:261-263; for *B. amyloliquefaciens* in Smith et al., (1986) Appl. and Env. Microbiol. 51:634; for *B. thuringiensis* in Fisher et al., (1981) Arch. Microbiol. 139:213-217; for *B. sphaericus* in McDonald (1984) J. Gen. Microbiol. 130:203; and *B. larvae* in Bakhiet et al., (1985, Appl. Environ. Microbiol. 49:577). Mann et al., (1986, Current Microbiol. 13:131-135) report on transformation of *Bacillus* protoplasts and Holubova, (1985) Folia Microbiol. 30:97) disclose methods for introducing DNA into protoplasts using DNA containing liposomes.

#### VI. Identification of Transformants

Whether a host cell has been transformed with a mutated or a naturally occurring gene encoding a gram-positive MP, detection of the presence/absence of marker gene expression can suggest whether the gene of interest is present. However, its expression should be confirmed. For example, if the nucleic acid encoding an MP of the present invention is inserted within a marker gene sequence, recombinant cells containing the insert can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with nucleic acid encoding the MP under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the MP as well.

Alternatively, host cells which contain the coding sequence for a metallo-protease and express the protein may be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridization and protein bioassay or immunoassay techniques which include membrane-based, solution-based, or chip-based technologies for the detection and/or quantification of the nucleic acid or protein.

The presence of the metallo-protease polynucleotide sequence can be detected by DNA-DNA or DNA-RNA hybridization or amplification using probes, portions or fragments of *B. subtilis* MP.

#### VII Assay of Protease Activity

There are various assays known to those of skill in the art for detecting and measuring protease activity. There are assays based upon the release of acid-soluble peptides from casein or hemoglobin measured as absorbance at 280 nm or colorimetrically using the Folin method

-- 16 --

(Bergmeyer, et al., 1984, Methods of Enzymatic Analysis vol. 5, Peptidases, Proteinases and their Inhibitors, Verlag Chemie, Weinheim). Other assays involve the solubilization of chromogenic substrates (Ward, 1983, Proteinases, in Microbial Enzymes and Biotechnology (W.M. Fogarty, ed.), Applied Science, London, pp. 251-317).

### VIII Secretion of Recombinant Proteins

Means for determining the levels of secretion of a heterologous or homologous protein in a gram-positive host cell and detecting secreted proteins include, using either polyclonal or monoclonal antibodies specific for the protein. Examples include enzyme-linked immunosorbent assay (ELISA), radioimmunoassay (RIA) and fluorescent activated cell sorting (FACS). These and other assays are described, among other places, in Hampton R et al (1990, Serological Methods, a Laboratory Manual, APS Press, St Paul MN) and Maddox DE et al (1983, J Exp Med 158:1211).

A wide variety of labels and conjugation techniques are known by those skilled in the art and can be used in various nucleic and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting specific polynucleotide sequences include oligolabeling, nick translation, end-labeling or PCR amplification using a labeled nucleotide. Alternatively, the nucleotide sequence, or any portion of it, may be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes *in vitro* by addition of an appropriate RNA polymerase such as T7, T3 or SP6 and labeled nucleotides.

A number of companies such as Pharmacia Biotech (Piscataway NJ), Promega (Madison WI), and US Biochemical Corp (Cleveland OH) supply commercial kits and protocols for these procedures. Suitable reporter molecules or labels include those radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents as well as substrates, cofactors, inhibitors, magnetic particles and the like. Patents teaching the use of such labels include US Patents 3,817,837; 3,850,752; 3,939,350; 3,996,345; 4,277,437; 4,275,149 and 4,366,241. Also, recombinant immunoglobulins may be produced as shown in US Patent No. 4,816,567 and incorporated herein by reference.

### IX Purification of Proteins

Gram positive host cells transformed with polynucleotide sequences encoding heterologous or homologous protein may be cultured under conditions suitable for the



-- 17 --

expression and recovery of the encoded protein from cell culture. The protein produced by a recombinant gram-positive host cell comprising a mutation or deletion of the metallo-protease activity will be secreted into the culture media. Other recombinant constructions may join the heterologous or homologous polynucleotide sequences to nucleotide sequence encoding a polypeptide domain which will facilitate purification of soluble proteins (Kroll DJ et al (1993) DNA Cell Biol 12:441-53).

Such purification facilitating domains include, but are not limited to, metal chelating peptides such as histidine-tryptophan modules that allow purification on immobilized metals (Porath J (1992) Protein Expr Purif 3:263-281), protein A domains that allow purification on immobilized immunoglobulin, and the domain utilized in the FLAGS extension/affinity purification system (Immunex Corp, Seattle WA). The inclusion of a cleavable linker sequence such as Factor XA or enterokinase (Invitrogen, San Diego CA) between the purification domain and the heterologous protein can be used to facilitate purification.

## X Uses of The Present Invention

### MP and Genetically Engineered Host Cells

The present invention provides genetically engineered host cells comprising mutations, preferably non-revertable mutations, or deletions in the naturally occurring gene encoding MP such that the proteolytic activity is diminished or deleted altogether. The host cell may contain additional protease deletions, such as deletions of the mature subtilisin protease and/or mature neutral protease disclosed in United States Patent No. 5,264,366.

In a preferred embodiment, the host cell is further genetically engineered to produce a desired protein or polypeptide. In a preferred embodiment the host cell is a *Bacillus*. In another preferred embodiment, the host cell is a *Bacillus subtilis*.

In an alternative embodiment, a host cell is genetically engineered to produce a gram-positive MP. In a preferred embodiment, the host cell is grown under large scale fermentation conditions. In another preferred embodiment, the MP is isolated and/or purified and used in the textile industry, the feed industry and in cleaning compositions such as detergents.

As noted, MP can be useful in formulating various cleaning compositions. A number of known compounds are suitable surfactants useful in compositions comprising the MP of the invention. These include nonionic, anionic, cationic, anionic or zwitterionic detergents, as disclosed in US 4,404,128 and US 4,261,868. A suitable detergent formulation is that described in Example 7 of US Patent 5,204,015. The art is familiar with the different

-- 18 --

formulations which can be used as cleaning compositions. In addition, MP can be used, for example, in bar or liquid soap applications, dishcare formulations, contact lens cleaning solutions or products, peptide hydrolysis, waste treatment, textile applications, as fusion-cleavage enzymes in protein production, etc. MP may comprise enhanced performance in a detergent composition (as compared to another detergent protease). As used herein, enhanced performance in a detergent is defined as increasing cleaning of certain enzyme sensitive stains such as grass or blood, as determined by usual evaluation after a standard wash cycle.

MP can be formulated into known powdered and liquid detergents having pH between 6.5 and 12.0 at levels of about .01 to about 5% (preferably .1% to .5%) by weight. These detergent cleaning compositions can also include other enzymes such as known proteases, amylases, cellulases, lipases or endoglycosidases, as well as builders and stabilizers.

The addition of MP to conventional cleaning compositions does not create any special use limitation. In other words, any temperature and pH suitable for the detergent is also suitable for the present compositions as long as the pH is within the above range, and the temperature is below the described MP's denaturing temperature. In addition, MP can be used in a cleaning composition without detergents, again either alone or in combination with builders and stabilizers.

Proteases can be included in animal feed such as part of animal feed additives as described in, for example, US 5,612,055; US 5,314,692; and US 5,147,642.

One aspect of the invention is a composition for the treatment of a textile that includes MP. The composition can be used to treat for example silk or wool as described in publications such as RD 216,034; EP 134,267; US 4,533,359; and EP 344,259.

#### MP Polynucleotides

A *B.subtilis* MP polynucleotide, or any part thereof, provides the basis for detecting the presence of gram-positive microorganism MP polynucleotide homologs through hybridization techniques and PCR technology.

Accordingly, one aspect of the present invention is to provide for nucleic acid hybridization and PCR probes which can be used to detect polynucleotide sequences, including genomic and cDNA sequences, encoding gram-positive MP or portions thereof. In another aspect of the present invention, an MP polynucleotide can be used in hybridization technology to detect the major protease of a gram-positive microorganism due to the proximity of the MP with the major protease.

-- 19 --

The manner and method of carrying out the present invention may be more fully understood by those of skill in the art by reference to the following examples, which examples are not intended in any manner to limit the scope of the present invention or of the claims directed thereto

### Example I

#### Preparation of a Genomic library

The following example illustrates the preparation of a *Bacillus* genomic library.

Genomic DNA from *Bacillus* cells is prepared as taught in Current Protocols In Molecular Biology vol. 1, edited by Ausubel et al., John Wiley & Sons, Inc. 1987, chapter 2. 4.1. Generally, *Bacillus* cells from a saturated liquid culture are lysed and the proteins removed by digestion with proteinase K. Cell wall debris, polysaccharides, and remaining proteins are removed by selective precipitation with CTAB, and high molecular weight genomic DNA is recovered from the resulting supernatant by isopropanol precipitation. If exceptionally clean genomic DNA is desired, an additional step of purifying the *Bacillus* genomic DNA on a cesium chloride gradient is added.

After obtaining purified genomic DNA, the DNA is subjected to Sau3A digestion. Sau3A recognizes the 4 base pair site GATC and generates fragments compatible with several convenient phage lambda and cosmid vectors. The DNA is subjected to partial digestion to increase the chance of obtaining random fragments.

The partially digested *Bacillus* genomic DNA is subjected to size fractionation on a 1% agarose gel prior to cloning into a vector. Alternatively, size fractionation on a sucrose gradient can be used. The genomic DNA obtained from the size fractionation step is purified away from the agarose and ligated into a cloning vector appropriate for use in a host cell and transformed into the host cell.

### Example II

#### Detection of gram-positive microorganisms

The following example describes the detection of gram-positive microorganism MP.

DNA derived from a gram-positive microorganism is prepared according to the methods disclosed in Current Protocols in Molecular Biology, Chap. 2 or 3. The nucleic acid is subjected to hybridization and/or PCR amplification with a probe or primer derived from MP.

-- 20 --

The nucleic acid probe is labeled by combining 50 pmol of the nucleic acid and 250 mCi of [gamma <sup>32</sup>P] adenosine triphosphate (Amersham, Chicago IL) and T4 polynucleotide kinase (DuPont NEN<sup>®</sup>, Boston MA). The labeled probe is purified with Sephadex G-25 super fine resin column (Pharmacia). A portion containing 10<sup>7</sup> counts per minute of each is used in a typical membrane based hybridization analysis of nucleic acid sample of either genomic or cDNA origin.

The DNA sample which has been subjected to restriction endonuclease digestion is fractionated on a 0.7 percent agarose gel and transferred to nylon membranes (Nytran Plus, Schleicher & Schuell, Durham NH). Hybridization is carried out for 16 hours at 40 degrees C. To remove nonspecific signals, blots are sequentially washed at room temperature under increasingly stringent conditions up to 0.1 x saline sodium citrate and 0.5% sodium dodecyl sulfate. The blots are exposed to film for several hours, the film developed and hybridization patterns are compared visually to detect polynucleotide homologs of *B.subtilis* MP. The homologs are subjected to confirmatory nucleic acid sequencing. Methods for nucleic acid sequencing are well known in the art. Conventional enzymatic methods employ DNA polymerase Klenow fragment, SEQUENASE<sup>®</sup> (US Biochemical Corp, Cleveland, OH) or Taq polymerase to extend DNA chains from an oligonucleotide primer annealed to the DNA template of interest.

Various other examples and modifications of the foregoing description and examples will be apparent to a person skilled in the art after reading the disclosure without departing from the spirit and scope of the invention, and it is intended that all such examples or modifications be included within the scope of the appended claims. All publications and patents referenced herein are hereby incorporated by reference in their entirety.